

**Discovery and characterization of endosymbiotic flagellates from Australian termites****Linda Ly, Coralyn Turner, Peter O'Donoghue**

University of Queensland, Australia

Termites are effective decomposers of wood, many utilizing endosymbiotic flagellates in their hindguts to aid digestion. Few studies have been conducted on the protistan fauna of Australian termites. Samples collected from 65 colonies of lower termites (Rhinotermitidae, Kalotermitidae, Mastotermitidae and Termopsidae) in north-eastern Australia were examined by light and electron microscopy. A total of 44 flagellate species were found belonging to 4 parabasalid orders (Cristamonadida, Trichonymphida, Spirotrichonymphida, and Trichomonadida) and one preaxostylid order (Oxymonadida). Only 14 flagellate species had previously been described, the remaining 30 were putative novel species. Oxymonads were only found in kalotermitids; cristamonads in all termite families except rhinotermitids; and trichomonads, trichonymphids and spirotrichonymphids were most abundant in rhinotermitids. The variable host specificity of the flagellates may be associated with host diet: kalotermitids feed on sound wood, termopsids on decaying wood and rhinotermitids on the whole range. Electron microscopy revealed many starch-feeding flagellates contained hydrogenosomes while most particulate-feeding flagellates contained hypersymbiotic bacteria, although several flagellates possessed neither.

1964-2014  
 **50**  
 Celebrating 50 years of the ASP

**2014 ASP 50th Anniversary Conference**  
 30th June to 3rd July 2014  
 ANU Commons, Canberra






# Programme & Abstracts

  
**ELSEVIER**

  
**IJP**  
INTERNATIONAL JOURNAL FOR PARASITOLOGY

  
**IJP**  
Drugs and Drug Resistance

  
**IJP**  
Parasites And Wildlife

  
**mla**  
MEAT & LIVESTOCK AUSTRALIA

  
**BAYER**

  
**Virbac**  
ANIMAL HEALTH

  
**Elanco**

  
**COMPOUNDS**  
AUSTRALIA

  
**BioLabs**  
NEW ENGLAND Inc.